

Download Yearbook Of Fishery Statistics Fao 1996

Capture Production Fao Statistics

(Source:MPEDA,Kochi) * Production from Aquaculture farms In India, brackish water aquaculture sector is largely based on farming of *Penaeus monodon*. Other shrimp species, i.e., *Penaeus indicus*, *P. merguensis* and *P. semisulcatus*, are considered as potential ones and cultivation of *Penaeus vannamei* has been gaining momentum in the recent years. In India, a major shift in India's policy on ...The Chinook salmon / ? ? ? n ? k / (*Oncorhynchus tshawytscha*) is the largest species in the Pacific salmon genus *Oncorhynchus*. The common name refers to the Chinookan peoples. Other vernacular names for the species include king salmon, Quinnet salmon, spring salmon, chrome hog, and Tyee salmon. The scientific species name is based on the Russian common name *chavycha* (??????). The largehead hairtail (*Trichiurus lepturus*) or beltfish is a member of the cutlassfish family, *Trichiuridae*. This common to abundant species is found in tropical and temperate oceans throughout the world. The taxonomy is not fully resolved, and the Atlantic, East Pacific and Northwest Pacific populations are also known as Atlantic cutlassfish, Pacific cutlassfish and Japanese cutlassfish ...How many animals does a vegetarian save? by Harish. Posted on February 6, 2012; Last updated on March 16, 2015. A vegetarian spares the lives of a certain number of animals each time he or she chooses to forgo meat for vegetables, fruits, grains, legumes and nuts.